

DFTUZ 92/14
 IPNO-TH 92/83
 September 1992

SECOND ORDER FORMALISM FOR FERMIONS: ANOMALIES AND SPECIES DOUBLING ON THE LATTICE [†]

J. L. Cortés¹, J. Gamboa² and L. Velázquez¹

¹*Departamento de Física Teórica, Universidad de Zaragoza,
 50009 Zaragoza, Spain.*

²*División de Physique Théorique, Institut de Physique Nucleaire[‡],
 F-91406 Cedex, Orsay, France.*

Abstract. A new formulation for fermions on the lattice based on a discretization of a second order formalism is proposed. A comparison with the first order formalism in connection with the $U(1)$ anomaly and the doubling problem is presented. The new formulation allows to eliminate half of the degrees of freedom, which can be important in non-perturbative calculations, and it can open new possibilities to the formulation of chiral gauge theories.

PACS 11.15.Ha , 11.30.Rd

[†] Work partially supported by CICYT (Proyecto AEN 90-0030).

[‡] *Unite de Recherche des Universités Paris 11 et Paris 6 associe au CNRS*

Many applications of relativistic quantum field theory in particle physics involve as an important ingredient a non-perturbative treatment of a fermionic degree of freedom, for which only limited results have been obtained. In a formulation based on a discretization of spacetime, it is only after a considerable progress in computing capabilities that, simulations going beyond the quenched approximations have been possible, although still with serious limitations. This makes interesting any attempt to simplify the treatment of a fermionic field.

Although in the case of theories involving parity conserving interactions the problem can, in principle, be reduced to a question of computing time, in the case of parity violating interactions the problem is more fundamental [1]. In fact almost nothing is known on non-perturbative aspects in chiral gauge theories as compared with the case of vector like theories. The basic difference is that no regularization which is manifestly gauge invariant is known (perhaps it does not exist).

When a gauge theory with fermionic matter fields is formulated on a lattice, then one finds that the only way to translate the gauge invariant solution of the doubling problem used in the case of vector like theories to the chiral case requires to use a scalar field. But, in fact, a recent work [2] has shown that these formulations fail because it is not possible to avoid really the doubling problem and the theories become vector like.

The other alternative is to use directly the solution of the doubling problem in the parity conserving case losing gauge invariance at the level of the lattice action. It is an open problem to see whether a gauge invariant continuum limit can be recovered and if this gives a framework where non-perturbative calculations can be done in practice.

At this point it is natural to look for a different starting point for a formulation of a fermionic field which could give a new perspective to the problem of including non-perturbative effects in the parity conserving case, and which could also be applied to chiral gauge theories.

In a previous work [3], a second order formulation of fermions ^{*}, based on the identification of decoupled field components in the usual Dirac action, has been introduced. It

^{*} In fact the possibility to consider a second order lattice action for fermions has been considered previously [4]

eliminates half of the integration variables in the path integral formulation and it leads to a different approach to chiral gauge theories. It can be interpreted as an elaborated version of an attempt by Feynman and Gell-Mann [5] to understand the properties of the weak interactions as a consequence of using a formulation based on the use of two components spinors instead of Dirac spinors.

The aim of this work is to explore the possibility to define a lattice second order formulation, first in the case of parity conserving gauge interactions, and the possible extension to chiral gauge theories.

The starting point of the second order formulation [3] is the identification of the variables

$$\begin{aligned}\chi_L &= \frac{1}{\sqrt{m}}\psi_L, & \chi_R &= \sqrt{m}(\psi_R + \frac{\vec{D}_L}{m}\psi_L), \\ \bar{\chi}_R &= \frac{1}{\sqrt{m}}\bar{\psi}_R, & \bar{\chi}_L &= \sqrt{m}(\bar{\psi}_L - \bar{\psi}_R \frac{\overleftarrow{D}_R}{m}),\end{aligned}\tag{1}$$

where ψ_L, ψ_R are the two chiral components of a Dirac field, which allow to rewrite the standard (euclidean) gauge invariant Dirac action

$$S^{(1)} = \int d^D x \left[m \bar{\psi} \psi + \frac{1}{2} \bar{\psi} (\vec{D} - \overleftarrow{D}) \psi \right],\tag{2}$$

into the form

$$S^{(1)} = \int d^D x \left[m^2 \bar{\chi}_R \chi_L + \bar{\chi}_R (\overleftarrow{D}_R \vec{D}_L) \chi_L + \bar{\chi}_L \chi_R \right].\tag{3}$$

From this expression one identifies $\bar{\chi}_L, \chi_R$ as auxiliary non- propagating fields leading to a second order formulation

$$S^{(2)} = \int d^D x \left[m^2 \bar{\chi}_R \chi_L + \bar{\chi}_R (\overleftarrow{D}_R \vec{D}_L) \chi_L \right],\tag{4}$$

which reproduces at the perturbative level all the results of the usual Dirac formulation including the $U(1)$ anomaly [3].

In the chiral case a mass term which is essential for the derivation of the second order formulation is not present in the Dirac action, but it is possible, in principle, to start directly with the second order action $S^{(2)}$ with $m = 0$ and the parity non- invariance of the gauge interaction is reflected on the different action of the covariant derivative on χ_L and $\bar{\chi}_R$.

In order to explore further the possibility to formulate a theory with fermionic degrees of freedom with a second order action, the next step is to introduce explicitly a regularization. One way to do that, which can be taken as a starting point for a non-perturbative study, is based on the formulation of the theory on a discrete lattice of spacetime points (a hypercubic lattice for example).

A naive discretization of the usual Dirac action leads to the well known doubling problem (the lattice action describes 2^D fermionic degrees of freedom for each fermionic field). This problem, which technically can be related to the first order action [6-7], is connected with the $U(1)$ anomaly. Since the anomaly is contained in the second order formulation [3] one can anticipate that the doubling problem will also be present in a discretization of the second order action although perhaps in a different way.

In order to show that and to study the way to avoid the doubling problem in the second order lattice formulation, let us translate to a lattice formulation the steps going from the Dirac action to the second order action.

The starting point is the naive discretization of the free Dirac action

$$S_f^{(1)} = \sum_{\mathbf{n}} \left[m \bar{\psi}_{\mathbf{n}} \psi_{\mathbf{n}} + \frac{1}{2} \bar{\psi}_{\mathbf{n}} \left(\overrightarrow{\Delta}^{\pm} - \overleftarrow{\Delta}^{\pm} \right) \psi_{\mathbf{n}} \right], \quad (5)$$

which is just the first order action in (1), particularized to the free case ($A_{\mu} = 0$), when the integral is replaced by a sum over the hypercubic lattice. The points of the lattice are characterized by a D-vector $\mathbf{n}$ whose components are integers in lattice units (a), the Dirac field $\psi(x)$ is replaced by a Grassmann variables $\psi_{\mathbf{n}}$ at each lattice point and the derivative is replaced by the symmetrized finite difference

$$(\Delta_{\mu}^{\pm} \psi)_{\mathbf{n}} = \frac{\psi_{\mathbf{n}+\hat{\mu}} - \psi_{\mathbf{n}-\hat{\mu}}}{2}, \quad (6)$$

where $\mathbf{n} \pm \hat{\mu}$ is a vector with components

$$(\mathbf{n} \pm \hat{\mu})_{\mu} = \mathbf{n}_{\mu} \pm 1; \quad (\mathbf{n} \pm \hat{\mu})_{\nu} = \mathbf{n}_{\nu} \text{ for } \nu \neq \mu. \quad (7)$$

The doubling problem is identified at this level in the discretized free fermion propagator, which is the free Dirac propagator with the momentum replaced by

$$\hat{P}_{\mu} = \frac{1}{a} \sin(a p_{\mu}), \quad (8)$$

vanishing not only at $p_\mu = 0$, but also at $p_\mu = \frac{\pi}{a}$, as a consequence of the discretization of the derivative in (6).

Nothing prevents us to repeat step by step the identification of decoupled variables and the naive second order action on the lattice will be

$$S_f^{(2)} = \sum_{\mathbf{n}} \left[m^2 \bar{\chi}_R \chi_L + \bar{\chi}_R \overleftrightarrow{\Delta}_R^\pm \overleftrightarrow{\Delta}_L^\pm \chi_L \right], \quad (9)$$

where the dynamical variables $\bar{\chi}_R, \chi_L$, as well as the decoupled auxiliary fields, will be given by (4) with the derivatives replaced by the finite difference (6).

Although the lattice action is of second order, as in the case of a bosonic field, the doubling problem is present due to the symmetrized finite difference approach for the derivative. This had to be the case since the action $S_f^{(2)}$ is nothing but a reformulation of the naive action $S_f^{(1)}$.

The only difference with the discretization of the Klein-Gordon action of a free bosonic field without any doubling problem is that in this case the kinetic operator is given by $\phi^+ \overleftrightarrow{\Delta}_\mu^+ \overleftrightarrow{\Delta}_\mu^+ \phi$ with the finite difference

$$(\Delta_\mu^+ \phi)_{\mathbf{n}} = \phi_{\mathbf{n}+\hat{\mu}} - \phi_{\mathbf{n}}, \quad (10)$$

instead of the symmetrized version in (6).

From this comparison it is very easy to identify what are the terms to be added to the naive action $S_f^{(2)}$ in order to avoid the doubling problem. By using the discretized version of the identity

$$\not{\partial}_R \not{\partial}_L = \partial_\mu \partial_\mu, \quad (11)$$

which is a consequence of the Dirac algebra and the commutation relation of the free derivatives, one is lead to consider

$$\bar{S}_f^{(2)} = S_f^{(2)} + \sum_{\mathbf{n}} \bar{\chi}_R \left(\overleftrightarrow{\Delta}_\mu^+ \overleftrightarrow{\Delta}_\mu^+ - \overleftrightarrow{\Delta}_\mu^\pm \overleftrightarrow{\Delta}_\mu^\pm \right) \chi_L, \quad (12)$$

as the free second order action.

The second term guarantees that the free propagator will have the same dependence in momentum as the bosonic propagator on the lattice and then it eliminates the doubling

problem. It plays the same role as the Wilson term in the first order action, being an irrelevant term in the naive classical continuum limit and giving a mass of the order of the cutoff to all the additional degrees of freedom which are lattice artifacts ^{*}.

In the presence of the gauge field one has the corresponding discretized versions of the covariant derivative

$$(D_\mu^\pm \psi)_\mathbf{n} = \frac{U_{\mathbf{n},\mu} \psi_{\mathbf{n}+\hat{\mu}} - U_{\mathbf{n},-\mu} \psi_{\mathbf{n}-\hat{\mu}}}{2}, \quad (13)$$

$$(D_\mu^+ \psi)_\mathbf{n} = U_{\mathbf{n},\mu} \psi_{\mathbf{n}+\hat{\mu}} - \psi_\mathbf{n}, \quad (14)$$

where $U_{\mathbf{n},\mu}$ is the gauge variable defined on the link between the points $\mathbf{n}$ and $\mathbf{n} + \hat{\mu}$. Covariant derivatives have a non trivial commutator and then one has

$$\overleftarrow{D}_R \overrightarrow{D}_L = \overleftarrow{D}_\mu \overrightarrow{D}_\mu + \frac{1}{4} [\gamma_\mu, \gamma_\nu]_{RL} (\overleftarrow{D}_\mu \overrightarrow{D}_\nu - \overleftarrow{D}_\nu \overrightarrow{D}_\mu), \quad (15)$$

where the first term is the covariant version of the free kinetic operator and the second term is due to the gauge interaction.

In order to identify the second order lattice formulation of a gauge theory what one has to do is to combine the free lattice action $\bar{S}_f^{(2)}$, when the derivatives Δ_μ are replaced by covariant derivatives, with an appropriate discretization of the contribution involving the commutator of covariant derivatives.

The simplest action one finds following this procedure is

$$\begin{aligned} S_{lat}^{(2)} = & \sum_{\mathbf{n}} \left[m^2 \bar{\chi}_R \chi_L + \bar{\chi}_R \overleftarrow{D}_\mu^+ \overrightarrow{D}_\mu^+ \chi_L \right] + \\ & + \frac{1}{4} \sum_{\mathbf{n}} \bar{\chi}_R [\gamma_\mu, \gamma_\nu]_{RL} (\overleftarrow{D}_\mu^\pm \overrightarrow{D}_\nu^\pm - \overleftarrow{D}_\nu^\pm \overrightarrow{D}_\mu^\pm) \chi_L, \end{aligned} \quad (16)$$

which corresponds to the minimal modification of the naive discretization of the Dirac action required in order to avoid the doubling problem. The term involving the commutator

^{*} In fact the operator in the second term

$$\overleftarrow{\Delta}_\mu^+ \overrightarrow{\Delta}_\mu^+ - \overleftarrow{\Delta}_\mu^\pm \overrightarrow{\Delta}_\mu^\pm = \left(\frac{\overleftarrow{\Delta}_\mu^+ - \overleftarrow{\Delta}_\mu^-}{2} \right) \left(\frac{\overrightarrow{\Delta}_\mu^+ - \overrightarrow{\Delta}_\mu^-}{2} \right),$$

is just the square of the operator which appears in the Wilson term.

of covariant derivatives does not require any modification and it is taken from the naive Dirac action which involves the symmetrized discrete version of the covariant derivative (13).

The action $S_{lat}^{(2)}$ is obtained from the naive discretization of the Dirac action by adding the term

$$\Delta S_{lat} = \sum_{\mathbf{n}} \bar{\chi}_R \left(\overleftarrow{D}_\mu^+ \overrightarrow{D}_\mu^+ - \overleftarrow{D}_\mu^\pm \overrightarrow{D}_\mu^\pm \right) \chi_L, \quad (17)$$

which is the gauge invariant version of the term identified in the free case as a way to solve the doubling problem.

The action $S_{lat}^{(2)}$ can be taken as a spacetime lattice regularization of the second order formulation based on the action $S^{(2)}$. As it is always the case in a lattice regularization, manifest covariance is lost, and all one has is an invariance under the discrete transformations leaving invariant the spacetime lattice ($\frac{\pi}{2}$ rotations in the hypercubic lattice) which one expects is enough to recover a covariant continuum limit.

The general analysis of the anomaly in the second order formulation [3] can be applied also to the lattice regularization. It was based on the identification of a regularization independent part and a combination of covariance arguments, to fix the more general tensor structure in the quadratic contribution to the effective action, together with the constraints imposed by gauge invariance. It is very easy to see that the invariance under lattice rotations is enough to fix the same tensor structure. Then a lattice regularization which is manifestly gauge invariant and which avoids the doubling problem, as it is the case for $S_{lat}^{(2)}$, allows to reproduce the standard result for the $U(1)$ anomaly. A part of the anomaly comes from the non invariance of the naive discretization of the second order action under a chiral transformation of the dynamical degrees of freedom $\chi_L, \bar{\chi}_R$, and the rest of the anomaly comes from the fermionic expectation value of the variation of the term ΔS_{lat} which is required in order to solve the doubling problem.

Once more one recognizes in the second order formulation the connection between the doubling problem and the anomaly. The only difference with the analysis based on the Dirac action is that the elimination of the decoupled fields $\chi_R, \bar{\chi}_L$ already introduces a violation of chiral invariance before the regularization is introduced.

The action $S_{lat}^{(2)}$ is just one of the possible ways to discretize the second order action $S^{(2)}$. It is natural to expect such an ambiguity for any regularization. The only properties that the correction ΔS_{lat} eliminating the doublers of the naive discretization should satisfy in order to reproduce the anomaly are gauge invariance and lattice rotational invariance. Then, unless unitarity imposes some additional constraints (we have not found a practical method to study this question), any lattice regularization free of doubling which differs from $S_{lat}^{(2)}$ in gauge invariant and lattice rotational invariant terms which are irrelevant in the naive continuum limit, can be used as a lattice second order formulation of a gauge theory.

At the perturbative level, a lattice second order formulation is equivalent to the standard first order formulation. From a practical point of view, as a consequence of the identification of the decoupled auxiliary fields $\chi_R, \bar{\chi}_L$, one has a path integral formulation with half of the anticommuting variables to integrate for a given lattice which, in principle, can be associated to an important “saving” of computing time.

This point, together with some other possible differences at the non-perturbative level, deserve further investigation. In particular the approach to the chiral massless limit could be one of the problems where the second order formulation could bring some new perspectives as a consequence of the different realization of chiral symmetry.

A related and still more promising possible application of a second order lattice formulation is to try to extend the action $S_{lat}^{(2)}$ to a chiral gauge theory. In fact a non-perturbative formulation of a chiral gauge theory is still an open problem; the attempts to extend a lattice first order formulation to the chiral case presents some basic difficulties which still do not have a satisfactory solution. This makes interesting any possibility to approach this problem from a different point of view, like the second order formulation.

One way to approach this problem, which is similar to the approach followed in the first order formulation [1], is based on the introduction of a scalar field coupled to the fermion field in order to give a mass to the fermion in a gauge invariant way. One can also apply this coupling to give masses to the replica fermions of the order of the cutoff in the chiral case through a gauge invariant Wilson-Yukawa term in the lattice action.

The translation of this approach to the second order formulation takes as starting

point an action

$$S_{ch}^{(1)} = \int d^D x \left[\bar{\psi}_R \phi \psi_L + \bar{\psi}_L \phi^* \psi_R + \frac{1}{2} \bar{\psi} (\vec{D} - \overleftarrow{D}) \psi \right], \quad (18)$$

where the mass term is replaced by a coupling to the scalar field

$$\begin{aligned} \chi_R &= \psi_R + \frac{1}{\phi^*} \vec{D}_L \psi_L, \\ \bar{\chi}_L &= \bar{\psi}_L - \bar{\psi}_R \overleftarrow{D}_R \frac{1}{\phi^*}, \end{aligned} \quad (19)$$

and then one has

$$S_{ch}^{(1)} = \int d^D x \left[\bar{\psi}_R \phi \psi_L + \bar{\psi}_L \overleftarrow{D}_R \frac{1}{\phi^*} \vec{D}_L \psi_L + \bar{\chi}_L \phi^* \chi_R \right], \quad (20)$$

The only term involving the variable χ_R is a trivial factor which can be absorbed into a redefinition of the fermionic measure, which leads to

$$S_{ch}^{(2)} = \int d^D x \left[\bar{\psi}_R \phi \psi_L + \bar{\psi}_L \overleftarrow{D}_R \frac{1}{\phi^*} \vec{D}_L \psi_L \right]. \quad (21)$$

It is not clear how to proceed from this non-polynomic action. If one formally translates the lattice regularization of the parity conserving case to the chiral action then one would have

$$\begin{aligned} S_{ch}^{(2)} &= \sum_{\mathbf{n}} \left[\bar{\psi}_R \phi \psi_L + \bar{\psi}_R \overleftarrow{D}_\mu^+ \frac{1}{\phi^*} \vec{D}_\mu^+ \psi_L \right] + \\ &+ \frac{1}{4} \sum_{\mathbf{n}} \bar{\psi}_R [\gamma_\mu, \gamma_\nu]_{RL} \left(\overleftarrow{D}_\mu^\pm \frac{1}{\phi^*} \vec{D}_\nu^\pm - \overleftarrow{D}_\nu^\pm \frac{1}{\phi^*} \vec{D}_\mu^\pm \right) \psi_L, \end{aligned} \quad (22)$$

which is the translation to the second order formalism of the formulation of a chiral theory based on the introduction of a scalar field. It is difficult to imagine why the difficulties which appear in the first order formalism [2] should not be present also in this case.

Another approach to the problem of regularizing a second order formulation of a chiral gauge theory is based on a direct discretization of the action (4) with $m = 0$ and the covariant derivative acting differently on $\bar{\chi}_R, \chi_L$. In this case one has the gauge variables $U_{\mathbf{n},\mu}$ and $\bar{U}_{\mathbf{n},\mu}$ which correspond to the group element in the representation of χ_L and $\bar{\chi}_R$ respectively.

In the abelian case one has

$$U_{\mathbf{n},\mu} = \exp(ieA_{\mathbf{n},\mu}); \quad \bar{U}_{\mathbf{n},\mu} = \exp(-i\bar{e}A_{\mathbf{n},\mu}). \quad (23)$$

The symmetrized discretized covariant derivatives will be

$$(\vec{D}_\mu^\pm \chi_L)_{\mathbf{n}} = \frac{U_{\mathbf{n},\mu} \chi_{L,\mathbf{n}+\hat{\mu}} - U_{\mathbf{n},-\mu} \chi_{L,\mathbf{n}-\hat{\mu}}}{2}, \quad (24a)$$

$$(\bar{\chi}_R \overleftarrow{D}_\mu^\pm)_{\mathbf{n}} = \frac{\bar{\chi}_{R,\mathbf{n}+\hat{\mu}} \bar{U}_{\mathbf{n},\mu} - \bar{\chi}_{R,\mathbf{n}-\hat{\mu}} \bar{U}_{\mathbf{n},-\mu}}{2} \quad (24b)$$

and

$$(\vec{D}_\mu^+ \chi_L)_{\mathbf{n}} = U_{\mathbf{n},\mu} \chi_{L,\mathbf{n}+\hat{\mu}} - \chi_{L,\mathbf{n}}, \quad (25a)$$

$$(\bar{\chi}_R \overleftarrow{D}_\mu^+)_{\mathbf{n}} = \bar{\chi}_{R,\mathbf{n}+\hat{\mu}} \bar{U}_{\mathbf{n},\mu} - \bar{\chi}_{R,\mathbf{n}}. \quad (25b)$$

Then one can translate the second order lattice formulation in the parity conserving case to the chiral case and one has

$$\begin{aligned} \bar{S}_{ch}^{(2)} = & \sum_{\mathbf{n}} \bar{\chi}_R \overleftarrow{D}_\mu^+ \vec{D}_\mu^+ \chi_L + \\ & + \frac{1}{4} \sum_{\mathbf{n}} \bar{\chi}_R [\gamma_\mu, \gamma_\nu]_{RL} (\overleftarrow{D}_\mu^\pm \vec{D}_\nu^\pm - \overleftarrow{D}_\nu^\pm \vec{D}_\mu^\pm) \chi_L, \end{aligned} \quad (26)$$

as a candidate for a lattice formulation of a chiral gauge theory.

The possibility to define a continuum limit which corresponds to a chiral gauge invariant relativistic quantum field theory from this lattice action remains as an open question. Note that the lattice action is not gauge invariant because it is already a regularization of a gauge non invariant second order action, and the additional term required to eliminate the doubling problem violates also gauge invariance. One can expect that the conservation of the current coupled to the gauge field in the case of an anomaly free fermion content will allow to find a gauge invariant continuum limit after appropriate counterterms are added. This seems to be a necessary step in order to get a chiral gauge invariant theory also in the approach based on the first order formalism.

Another non trivial question is to see whether the continuum limit corresponds to a unitary theory, also as a consequence of the conservation of the current coupled to the

gauge field. We do not know how to prove the validity of the formulation of a chiral gauge theory based on the lattice action $\bar{S}_{ch}^{(2)}$. As a first step, a perturbative and non-perturbative analysis of two dimensional models is under investigation.

Other open problems, like the introduction of four fermion interactions in the second order formalism, the study of spontaneous symmetry breaking and dynamical generation of mass, will be the subject of a future work.

To summarize a second order formulation based on the identification of a combination of fermionic field components with no dynamics has been proposed as a way to study a gauge invariant parity conserving theory. The possibility to apply this formalism to the case of chiral gauge theories has been also pointed out with the new perspectives that it can open on the dynamics of these theories.

A lattice regularization of the second order parity invariant formulation free of the doubling problem has been presented. The identification of the decoupled auxiliary fields leads to a path integral formulation with half of the integration variables which can simplify considerably any numerical study.

An attempt to give a non-perturbative formulation of chiral gauge theories on the lattice has been presented in detail. It remains for the future to test the validity of this formulation and to apply it to study some non-perturbative effects of the standard model.

We would like to thank H.B. Nielsen by useful discussions.

References

- [1] For a review and references see M.F.L.Golterman, Nucl.Phys. **20B** (Proc.Suppl.) (1991)528.
- [2] M.F.L.Golterman, D.N.Petcher and J.Smit, Nucl.Phys. **370B**(1992)51.

- [3] J.L. Cortés, J. Gamboa and L.Velázquez, *Second Order Formalism for Fermions* Zaragoza-Orsay Preprint, DFTUZ 92/13.
- [4] T. Banks and A. Casher, Nucl. Phys. **169B**(1980)103; A.C. Longhitano and B. Svetitsky, Phys Lett. **126B**(1983)259.
- [5] R.P. Feynman and M. Gell-Mann, Phys. Rev. **109**(1958)193.
- [6] K.G. Wilson e.g *New Phenomena in Subnuclear Physics*, ed. A. Zichichi, Plenum (1977).
- [7] H.B. Nielsen and M. Ninomiya, Nucl. Phys. **185B**(1981)20; **193B**(1981)173; Phys. Lett. **105B**(1981)219.